




Huma Gro[®] International Report

Justin Smith- Vice President Huma Gro® International



Percentage of Domestic and International Sales of Bio Huma Netics, Inc., 2004 - 2013



HUMA C-AGRO



Ten Languages Spoken by BHN Employees

- English -40
- Spanish 9
- Portuguese 4
- French 3
- Chinese 2
- German 2
- Arabic 1
- Finnish 1
- Japanese- 1
- Hungarian 1
- Italian 1

2014 Huma Gro[®] Worldwide Presence





Finding New Huma Gro Distributors

• Trade Shows/Expos

-Argus/FMB-Colombia

-FCI

-New Ag Intl-Poland

-CAC-Slovenia

-World Ag Expo-Vietnam, El Salvador

- US Commercial Services (Department of Commerce)
 -Gold Key Service, Advertisements-Spain
- Website, Social Media, Marketing, Magazines-Israel, Philippines



HUMA CAGRO

Steps to Opening a New Country

- Initial Contact, Mutual Interest
- Face to face Meetings
 - Dist training, grower visits, infield work
- Send Samples of Key Products
- Contract
- Begin Registration Process (if necessary)
- Begin Field Demonstrations of Products
- Products Registered
- First Container

Ordered









The HUMA GRO[®] Phosphorus Fertilizer Label SUPER PHOS[®]- In Many Markets

SUPER PHOS TA CLARANTEED ANALYSIS 0-50-0 Available Progebase (P,O) Deniesed from: Progebase: Add Deniesed from: Progebase: Add				1 每国优马(BHN)公司 会球20多个国家或地区, 0多年的一种专业助利 品采用"物理技术"。 有利题片段结合数体, 主成分及天然阔,能推 疗或药物分子,有效提 "药的时用罪。"	tt.
METHOD OF APPLICATION	SUGGESTED RATE PER HECTARE				
METHOD OF APPLICATION	Field Crops	Tree or Vine Crops		Line & Statement and all der tem	
Foliar band application at 50% coverage	up to 5 liters			!混合使用,减少氨肥	
Foliar broadcast or sprinklers: solid, set, pivot, linear (100% speed)	up to 10 liters	up to 20 liters		: 50%:	
Soil banded/ injected/ drip tape/ micro sprinklers	up to 20 liters	up to 30 liters		閉肥料 (复合肥或复混肥)	
Soil broadcast or furrow irrigated	up to 30 liters	up to 40 liters		,减少用量达30%:	
DIRECTODS 	SUPER PHOS ^{IM} should be applied cation is concentrated in the active lizers. If compatibility is in questi to not apply this product foliarly in n for follar application. I bo not app d rates may vary according to indi tions can be made as often as eve endations. Do not store this pro	root zone or on the leaf surface. on, jar test a small quantity. App concentrations greater than 5%. by HUMA GRO [®] nutrients during in vidual crop requirement, stage o ry 7 to 10 days as needed. «Co duct below 10°C (50°F) or above	 Can be applied ply foliar sprays For maximum the hottest time if development, nsult your local 30°C (90°F). 	的流失造成肥料浪费 肥,可长期保持肥料 和稳定性; 高农药的药效。 そ	
Control The second se	RO by	151 Kilogenen per Line (J. 197 C. 127 Sa. per grade (G. 197 F. per 4.3. 128 Sa. per sales (G. 197 F. 129 Sa. per sales (G. 197 Sa. 198 Sa. 199 Sa.	NG-105-0152012	E肥料颗粒的表面,混)后,尿素、碳铵施肥 -半,施肥方法按原来 :法施肥。每 [@] (每 50 公 150ml 减伴 [®])。	ŧ

English



Mandarin Chinese



Turkish





International Travel – an Important Element of BHN's Success



Guatemala




Nicaragua




Mexico



Turkey



China



Taiwan



THANK YOU!